

FLOOD & DROUGHT BULLETIN

APRIL 2026




Table of Contents

| | |
|--|----|
| INTRODUCTION | 3 |
| THE STANDARDIZED PRECIPITATION INDEX (SPI) | 3 |
| 1-month SPI..... | 4 |
| 3-month SPI..... | 4 |
| 6-month SPI..... | 4 |
| 12-month SPI..... | 4 |
| SPI ANALYSIS FOR APRIL 2026..... | 6 |
| 1-Month SPI (April 2026)..... | 6 |
| 3-month SPI (February 2026 - April 2026)..... | 6 |
| 6-month SPI (November 2025 - April 2026) | 7 |
| 12-month SPI (May 2025 - April 2026)..... | 7 |
| STATIONS | 9 |
| REFERENCE..... | 10 |

INTRODUCTION

The Flood and Drought bulletin is a monthly analysis of rainfall in Ghana prepared and released by the Ghana Meteorological Agency (GMet). The bulletin provides an authoritative and scientific understanding of rainfall variability which is essential for accurate hydrological modeling, climate change assessments, and effective water resource planning to mitigate risks associated with extreme events like droughts and floods.

Among other services and products, the Flood and Drought bulletin complements the objectives of GMet in line with the National Framework for Climate Services (NFCS) to improve co-production, tailoring, delivery and use of science-based climate predictions and services focused on the five pillars of the Global Framework for Climate Services (GFCS) by the World Meteorological Organization (WMO): agriculture and food security, disaster risk reduction, energy, health and water.

The analysis in the Flood and Drought bulletin is based on the Standardized Precipitation Index (SPI) developed by McKee et al. (1993) for the purpose of *defining and monitoring drought*. Drought is an insidious natural hazard that results from lower levels of precipitation than what is considered normal. When this phenomenon extends over a season or a longer period of time, water becomes increasingly insufficient to meet the demands of human activities and the environment. Drought must be considered a relative, rather than absolute, condition. Drought means different things to different users such as water managers, agricultural producers, hydroelectric power plant operators and wildlife biologists. Even within sectors, there are many different perspectives of drought because impacts may differ markedly. Droughts are commonly classified by type as meteorological, agricultural and hydrological, and differ from one another in intensity, duration and spatial coverage (WMO, 2012).

THE STANDARDIZED PRECIPITATION INDEX (SPI)

SPI indicator, which was developed by McKee et al. (1993), and described in detail by Edwards and McKee (1997), measures precipitation anomalies at a given location, based on a comparison of observed total precipitation amounts for an accumulation period of interest (e.g. 1, 3, 12, 48 months), with the long-term historic rainfall record for that period. This indicator measures anomalies of accumulated precipitation during a given period. In calculating SPI, precipitation is the only required input parameter (McKee and others, 1993, 1995). The SPI calculation for any location is based on the long-term precipitation record for a desired period. This long-term record is fitted to a probability distribution, which is then transformed into a normal distribution so that the mean SPI for the location and desired period is zero (Edwards and McKee, 1997).

The SPI is designed to quantify the precipitation deficit for multiple timescales and can be calculated from 1 month up to 72 months. Statistically, 1–24 months is the best practical range of

application (Guttman, 1994, 1999). The Flood and Drought bulletin of Ghana considers only the 1-month, 3-month, 6-month and 12-month SPI.

1-month SPI

The 1-month SPI compares the precipitation of a specific month with the precipitation totals from the same month for all the years included in the historical record.

3-month SPI

The 3-month SPI provides a comparison of the precipitation over a specific 3 consecutive month period with the precipitation totals from the same 3-month period for all the years included in the historical record.

6-month SPI

The 6-month SPI compares the precipitation for a specific 6 consecutive months with the same 6-month period over the historical record.

12-month SPI

The 12-month SPI is a comparison of the precipitation for 12 consecutive months with that recorded in the same 12 months in all previous years of available data.

These timescales reflect the impact of drought on the availability of the different water resources. Meteorological and soil moisture conditions (agriculture) respond to precipitation anomalies on relatively short timescales, for example 1 to 3 months (SPI-1 to SPI-3), whereas streamflow, reservoirs, and groundwater respond to longer-term precipitation anomalies, for example 3 months to 12 months (SPI-3 to SPI-12) or longer (EDO, 2020).

A real strength of the SPI is its ability to be calculated for many timescales, which makes it possible to deal with many of the drought types described above. The ability to compute the SPI on multiple timescales allows for temporal flexibility in the evaluation of precipitation conditions in relation to water supply.

In the Flood and Drought bulletin, the SPI values for any given location and accumulation period, are classified into nine different precipitation regimes (from dry to wet), as shown in Table 1. As can be seen, increasingly severe rainfall deficits (i.e., meteorological droughts) are indicated as SPI decreases below -0.5, while increasingly severe excess rainfall is indicated as SPI increases above 0.5.










| <i>ANOMALY</i> | <i>RANGE OF SPI VALUES</i> | <i>PRECIPITATION REGIME</i> | <i>COLOUR</i> |
|----------------|----------------------------|-----------------------------|---|
| Negative | Min \leq SPI \leq -2.0 | Extreme dry |  |
| | -2.0 < SPI \leq -1.5 | Severe dry |  |
| | -1.5 < SPI \leq -1.0 | Moderate dry |  |
| | -1.0 < SPI \leq -0.5 | Mild dry |  |
| None | -0.5 < SPI \leq 0.5 | Normal precipitation |  |
| Positive | 0.5 < SPI \leq 1.0 | Mild wet |  |
| | 1.0 < SPI \leq 1.5 | Moderate wet |  |
| | 1.5 < SPI \leq 2.0 | Severe wet |  |
| | 2.0 < SPI \leq Max | Extreme wet |  |

Table 1: SPI classification scheme used in the Flood and Drought bulletin

SPI ANALYSIS FOR APRIL 2026

The SPI analysis for rainfall in April 2026 has been generated in multiple timescales of 1-month (April 2026), 3-month (February 2026 – April 2026), 6-month (November 2025 – April 2026) and 12-month (May 2025 – April 2026). The maps generated depict the severity (positive or negative) of rainfall anomalies for the period under review.




Fig. 1(a): 1-Month SPI (for meteorological drought): April 2026




Fig. 1(b): 3-Month SPI (for agricultural drought): February 2026 – April 2026

1-Month SPI (April 2026)

The 1-Month SPI for April 2026 in Fig. 1(a) depicts predominantly mild dry conditions across the Savannah zone of the country, except for Walewale which showed near normal conditions, while areas such as Bole and Yendi depicted moderate dry conditions. Within the Transition zone, Salaga recorded moderate dryness whereas Wenchi, Kintampo and Ejura showed mild dry conditions. Sunyani, Dormaa, Prang and Atebubu depicted near normal to mild wet conditions, while Bechem and Kete Krachi displayed moderate wet conditions. The Forest zone showed largely mild to moderate wet conditions with Half Assini and Akim Oda recording severe wet conditions, whereas Abetifi depicted mild dryness. Along the Coastal zone, mild to moderate wet conditions were recorded in most areas, with Accra depicting near normal conditions.

3-month SPI (February 2026 - April 2026)

The 3-Month SPI, as shown in Fig. 1(b), indicates predominantly near-normal to mildly wet conditions across most parts of the country. In the Savannah zone, areas such as Navrongo, Bolgatanga, and Zuarungu experienced mild dryness, whereas Wa and Walewale recorded mild to moderate wet conditions. Bole, however, exhibited moderate dry conditions. In the Transition zone, mild wet conditions prevailed across most areas, including Kete Krachi, Prang, and Atebubu, while Kintampo, Wenchi, and Sunyani experienced moderate wetness. Salaga was the only area that recorded moderate dryness. Within the Forest zone, locations such as Axim, Takoradi, Sefwi Bekwai, Dunkwa, Kpando, Enchi, Kade, Asamankese, Koforidua, Akuse, and Ho experienced mild to moderate wet conditions. Bechem and Akim Oda recorded severe wetness, whereas Ejura exhibited near-normal conditions. The Coastal zone generally experienced mild to moderate wet conditions, with severe wetness observed in Cape Coast and Akatsi.




Fig. 1(c): 6-Month SPI (for hydrological drought): November 2025 – April 2026




Fig. 1(d): 12-Month SPI (for streamflow and lake storage drought): May 2025 – April 2026

6-month SPI (November 2025 - April 2026)

The 6-Month SPI, as shown in Fig. 1(c), indicates predominantly near-normal conditions across parts of the eastern Savannah zone. However, areas such as Navrongo, Bolgatanga, Zuarungu, and Yendi experienced mild to moderate dry conditions, while Walewale recorded moderate wetness. Bole and its environs also exhibited near normal to mild dry conditions while Wa experienced severe wetness. In the Transition zone, mild to moderate wet conditions prevailed in areas such as Kintampo, Wenchi, Prang, Atebubu, and Kete Krachi. Salaga, however, recorded moderate dryness, while Sunyani and Bechem experienced severe wet conditions. Within the Forest zone,

locations including Half Assini, Enchi, Axim, Goaso, Sefwi Bekwai, Ejura, Dunkwa, Kumasi, Akuse, Ho, and Kpando recorded mild to moderate wet conditions. Meanwhile, Abetifi, Akim Oda, Koforidua, and Asamankese experienced severe to extreme wetness. The Coastal zone generally exhibited moderate to severe wet conditions across most areas.

12-month SPI (May 2025 - April 2026)

The 12-Month SPI as shown in Fig. 1(d) indicates mild dry conditions dominating most parts of the Savannah zone including Zuarungu, Bolgatanga, Tamale and Yendi while Bole recorded moderate dryness. Walewale, Navrongo and Wa displayed mild to moderate wetness. Within the Transition zone, Salaga recorded severe to extreme dry conditions, with surrounding areas including Kete Krachi and Atebubu depicting moderate dry conditions while Kintampo, Dormaa and Prang recorded near normal conditions. Sunyani and Bechem, however, displayed moderate to severe wet conditions. The Forest zone in these areas; Kumasi, Goaso, Sefwi Bekwai, Dunkwa, Koforidua, Abetifi and Akuse recorded mild to moderate wet conditions, while Akim Oda, depicted severe wetness. Enchi and Dunkwa also recorded mild dryness. Along the Coastal zone, areas like, Cape Coast, Saltpond, Accra, Tema and Akatsi generally showed mild to moderate wet conditions.

| Station | Abbreviation | Station | Abbreviation | Station | Abbreviation |
|------------|--------------|--------------|--------------|----------|--------------|
| Abetifi | ABE | Enchi | ENC | Sunyani | SUN |
| Accra | ACC | Goaso | GOA | Takoradi | TDI |
| Akatsi | AKA | Half Assini | H ASSINI | Tamale | TLE |
| Akim Oda | ODA | Ho | HO | Tema | TEM |
| Akuse | AKU | Kade | KADE | Wa | WA |
| Asamankese | ASA | Kete Krachi | KRA | Walewale | WAL |
| Atebubu | ATE | Kintampo | KIN | Wenchi | WEN |
| Axim | AXM | Koforidua | KDA | Yendi | YDI |
| Bechem | BEC | Kpandu | KPA | Zuarungu | ZUA |
| Bole | BOL | Kumasi | KSI | | |
| Bolga | BOLGA | Mim | MIM | | |
| Bui | BUI | Navrongo | NAV | | |
| Cape Coast | C_COAST | Prang | PRANG | | |
| Dormaa | DOR | Salaga | SGA | | |
| Dunkwa | DUN | Saltpond | SAL | | |
| Ejura | EJU | Sefwi Bekwai | BEK | | |

STATIONS

REFERENCE

Copernicus European Drought Observatory (EDO): <https://edo.jrc.ec.europa.eu/> © European Commission, 2020.

Edwards, D. C. and T. B. McKee, 1997: Characteristics of 20th century drought in the United States at multiple time scales. Climatology Report 97-2, Department of Atmospheric Science, Colorado State University, Fort Collins, Colorado.

Guttman, N.B., 1994: On the sensitivity of sample L moments to sample size. *Journal of Climate*, 7(6):1026–1029.

———, 1999: Accepting the Standardized Precipitation Index: a calculation algorithm. *Journal of the American Water Resources Association*, 35(2):311–322.

McKee, T.B., N.J. Doesken and J. Kleist, 1993: The relationship of drought frequency and duration to time scale. In: *Proceedings of the Eighth Conference on Applied Climatology*, Anaheim, California, 17–22 January 1993. Boston, American Meteorological Society, 179–184.

———, 1995: Drought monitoring with multiple timescales. In: *Proceedings of the Ninth Conference on Applied Climatology*, Dallas, Texas, 15–20 January 1995. Boston American Meteorological Society, 233–236.

World Meteorological Organization, 2012: *Standardized Precipitation Index User Guide* (M. Svoboda, M. Hayes and D. Wood). (WMO-No. 1090), Geneva.

**Kindly send feedback to:
The Director-General
Ghana Meteorological Agency
P. O. Box Lg 87, Legon, Accra-Ghana
Email: info@meteo.gov.gh
Phone: 0302764926 / 0302777172**