


Table of Contents

| INTRODUCTION | |
|--|---|
| THE STANDARDIZED PRECIPITATION INDEX (SPI) | |
| 1-month SPI | 4 |
| 3-month SPI | 4 |
| 6-month SPI | 4 |
| 12-month SPI | 4 |
| SPI ANALYSIS FOR NOVEMBER 2025 | 6 |
| 1-Month SPI (November 2025) | 6 |
| 3-month SPI (September 2025 - November 2025) | 6 |
| 6-month SPI (June 2025 - November 2025) | 7 |
| 12-month SPI (December 2024 - November 2025) | 7 |
| STATIONS | 8 |
| REFERENCE | q |



0553538292



INTRODUCTION

The Flood and Drought bulletin is a monthly analysis of rainfall in Ghana prepared and released by the Ghana Meteorological Agency (GMet). The bulletin provides an authoritative and scientific understanding of rainfall variability which is essential for accurate hydrological modeling, climate change assessments, and effective water resource planning to mitigate risks associated with extreme events like droughts and floods.

Among other services and products, the Flood and Drought bulletin complements the objectives of GMet in line with the National Framework for Climate Services (NFCS) to improve coproduction, tailoring, delivery and use of science-based climate predictions and services focused on the five pillars of the Global Framework for Climate Services (GFCS) by the World Meteorological Organization (WMO): agriculture and food security, disaster risk reduction, energy, health and water.

The analysis in the Flood and Drought bulletin is based on the Standardized Precipitation Index (SPI) developed by McKee et al. (1993) for the purpose of *defining and monitoring drought*. Drought is an insidious natural hazard that results from lower levels of precipitation than what is considered normal. When this phenomenon extends over a season or a longer period, water becomes increasingly insufficient to meet the demands of human activities and the environment. Drought must be considered a relative, rather than absolute, condition. Drought means different things to different users such as water managers, agricultural producers, hydroelectric power plant operators and wildlife biologists. Even within sectors, there are many different perspectives of drought because impacts may differ markedly. Droughts are commonly classified by type as meteorological, agricultural and hydrological, and differ from one another in intensity, duration and spatial coverage (WMO, 2012).

THE STANDARDIZED PRECIPITATION INDEX (SPI)

SPI indicator, which was developed by McKee et al. (1993), and described in detail by Edwards and McKee (1997), measures precipitation anomalies at a given location, based on a comparison of observed total precipitation amounts for an accumulation period of interest (e.g. 1, 3, 12, 48 months), with the long-term historic rainfall record for that period. This indicator measures anomalies of accumulated precipitation during a given period. In calculating SPI, precipitation is the only required input parameter (McKee and others, 1993, 1995). The SPI calculation for any location is based on the long-term precipitation record for a desired period. This long-term record is fitted to a probability distribution, which is then transformed into a normal distribution so that the mean SPI for the location and desired period is zero (Edwards and McKee, 1997).

The SPI is designed to quantify the precipitation deficit for multiple timescales and can be calculated from 1 month up to 72 months. Statistically, 1–24 months is the best practical range of









application (Guttman, 1994, 1999). The Flood and Drought bulletin of Ghana considers only the 1-month, 3-month, 6-month and 12-month SPI.

1-month SPI

The 1-month SPI compares the precipitation of a specific month with the precipitation totals from the same month for all the years included in the historical record.

3-month SPI

The 3-month SPI provides a comparison of the precipitation over a specific 3 consecutive month period with the precipitation totals from the same 3-month period for all the years included in the historical record.

6-month SPI

The 6-month SPI compares the precipitation for a specific 6 consecutive months with the same 6-month period over the historical record.

12-month SPI


The 12-month SPI is a comparison of the precipitation for 12 consecutive months with that recorded in the same 12 months in all previous years of available data.

These timescales reflect the impact of drought on the availability of the different water resources. Meteorological and soil moisture conditions (agriculture) respond to precipitation anomalies on relatively short timescales, for example 1 to 3 months (SPI-1 to SPI-3), whereas streamflow, reservoirs, and groundwater respond to longer-term precipitation anomalies, for example 3 months to 12 months (SPI-3 to SPI-12) or longer (EDO, 2020).

A real strength of the SPI is its ability to be calculated for many timescales, which makes it possible to deal with many of the drought types described above. The ability to compute the SPI on multiple timescales allows for temporal flexibility in the evaluation of precipitation conditions in relation to water supply.

In the Flood and Drought bulletin, the SPI values for any given location and accumulation period, are classified into nine different precipitation regimes (from dry to wet), as shown in Table 1. As can be seen, increasingly severe rainfall deficits (i.e., meteorological droughts) are indicated as SPI decreases below -0.5, while increasingly severe excess rainfall is indicated as SPI increases above 0.5.








| | | PRECIPITATION | |
|----------|---------------------|----------------------|--------|
| ANOMALY | RANGE OF SPI VALUES | REGIME | COLOUR |
| Negative | Min <= SPI <= -2.0 | Extreme dry | |
| | -2.0 < SPI <= -1.5 | Severe dry | |
| | -1.5 < SPI <= -1.0 | Moderate dry | |
| | -1.0 < SPI <= -0.5 | Mild dry | |
| None | -0.5 < SPI <= 0.5 | Normal precipitation | |
| Positive | 0.5 < SPI <= 1.0 | Mild wet | |
| | 1.0 < SPI <= 1.5 | Moderate wet | |
| | 1.5 < SPI <= 2.0 | Severe wet | |
| | 2.0 < SPI <= Max | Extreme wet | |

Table 1: SPI classification scheme used in the Flood and Drought bulletin



0553538292



SPI ANALYSIS FOR NOVEMBER 2025

The SPI analysis for rainfall in September 2025 has been generated in multiple timescales of 1-month (November 2025), 3-month (September 2025 – November 2025), 6-month (June 2025 – November 2025) and 12-month (December 2024 – November 2025). The maps generated depict the severity (positive or negative) of rainfall anomalies for the period under review.




Fig. 1(a): 1-Month SPI (for meteorological drought): November 2025





Fig. 1(b): 3-Month SPI *(for agricultural drought)*: September 2025 – November 2025

1-Month SPI (November 2025)

Figure 1(a) presents the 1-Month Standardized Precipitation Index (SPI) for November 2025 is shown in Fig. 1. Mild to severe wet conditions are seen almost all over the country with spots of normal conditions around Atebubu, Koforidua, Akuse and Dormaa. Salaga observed mild dry conditions while it environs observed normal conditions.

3-month SPI (September 2025 - November 2025)

The 3-Month SPI shown in Fig. 1(b) reveals mild to moderate wet conditions over most parts of the country. Areas beneath the northern region through to the transition zone recorded normal conditions with areas such as Salaga, Ejisu and Bole recording mild to moderate dry conditions. Also, areas such Enchi, Dunkwa and Kade observed mild dry conditions.












Fig. 1(c): 6-Month SPI *(for hydrological drought)*: June 2025 – November 2025




Fig. 1(d): 12-Month SPI (for streamflow and lake storage drought): December 2024 – November 2025


6-month SPI (June 2025 - November 2025)

The 6-month SPI shown in Fig. 1(c) reveals normal to dry conditions in almost all over the country. Normal to moderately wet conditions were recorded around the coastal areas and parts of the forest zone (Abetifi, Kpando, Goaso and Mim) and transitions zone (Wenchi, Sunyani and Bechem). Also, most regions in the upper area of the country observed normal to mild wet conditions with Navrongo and Bolga recording mild dry conditions.

12-month SPI (December 2024 - November 2025)

The 12-Month SPI Fig. 1(d) for monitoring streamflow and lake storage indicates normal to dry conditions in almost all over the country. Normal to moderately wet conditions were recorded around the coastal areas and parts of the forest zone (Abetifi, Kpando, Goaso, Abetifi, Kpando, Ho and Mim). A deficit was observed around the upper east environs compared to the 6-months SPI.








STATIONS

| Station | Abbreviation | Station | Abbreviation | Station | Abbreviation |
|------------|--------------|--------------|--------------|----------|--------------|
| Abetifi | ABE | Enchi | ENC | Sunyani | SUN |
| Accra | ACC | Goaso | GOA | Takoradi | TDI |
| Akatsi | AKA | Half Assini | H_ASSINI | Tamale | TLE |
| Akim Oda | ODA | Но | НО | Tema | TEM |
| Akuse | AKU | Kade | KADE | Wa | WA |
| Asamankese | ASA | Kete Krachi | KRA | Walewale | WAL |
| Atebubu | ATE | Kintampo | KIN | Wenchi | WEN |
| Axim | AXM | Koforidua | KDA | Yendi | YDI |
| Bechem | BEC | Kpandu | KPA | Zuarungu | ZUA |
| Bole | BOL | Kumasi | KSI | | |
| Bolga | BOLGA | Mim | MIM | | |
| Bui | BUI | Navrongo | NAV | | |
| Cape Coast | C_COAST | Prang | PRANG | | |
| Dormaa | DOR | Salaga | SGA | | |
| Dunkwa | DUN | Saltpond | SAL | | |
| Ejura | EJU | Sefwi Bekwai | BEK | | |







REFERENCE

Copernicus European Drought Observatory (EDO): https://edo.jrc.ec.europa.eu/ © European Commission, 2020.

Edwards, D. C. and T. B. McKee, 1997: Characteristics of 20th century drought in the United States at multiple time scales. Climatology Report 97-2, Department of Atmospheric Science, Colorado State University, Fort Collins, Colorado.

Guttman, N.B., 1994: On the sensitivity of sample L moments to sample size. Journal of Climate, 7(6):1026–1029.


———, 1999: Accepting the Standardized Precipitation Index: a calculation algorithm. Journal of the American Water Resources Association, 35(2):311–322.

McKee, T.B., N.J. Doesken and J. Kleist, 1993: The relationship of drought frequency and duration to time scale. In: Proceedings of the Eighth Conference on Applied Climatology, Anaheim, California, 17–22 January 1993. Boston, American Meteorological Society, 179–184.

———, 1995: Drought monitoring with multiple timescales. In: Proceedings of the Ninth Conference on Applied Climatology, Dallas, Texas, 15–20 January 1995. Boston American Meteorological Society, 233–236.

World Meteorological Organization, 2012: *Standardized Precipitation Index User Guide* (M. Svoboda, M. Hayes and D. Wood). (WMO-No. 1090), Geneva.







Kindly send feedback to:
The Director-General
Ghana Meteorological Agency
P. O. Box Lg 87, Legon, Accra-Ghana
Email: info@meteo.gov.gh

Phone: 0302764926 / 0302777172